

Constraining a matter-dominated cosmological model with bulk viscosity proportional to the Hubble parameter

Arturo Avelino and Ulises Nucamendi

*Instituto de Física y Matemáticas
Universidad Michoacana de San Nicolás de Hidalgo
Edificio C-3, Ciudad Universitaria, CP. 58040
Morelia, Michoacán, México*

Abstract

We present and constrain a cosmological model where the only component is a pressureless fluid with bulk viscosity as an explanation for the present accelerated expansion of the universe. We study the particular model of a bulk viscosity coefficient proportional to the Hubble parameter: $\zeta_m = \zeta H$, where $\zeta = \text{constant}$. The possible values of ζ are constrained using the SNe Ia Gold 2006 sample, the CMB shift parameter R from the three-year WMAP observations, the Baryon Acoustic Oscillation (BAO) peak A from the Sloan Digital Sky Survey (SDSS) and the Second Law of Thermodynamics (SLT). It was found that this model is in agreement with the SLT using only the SNe Ia test. However when the model is constrained using the three cosmological tests together (SNe+CMB+BAO) the results are: 1.- The model violates the SLT, 2.- It predicts a value of $H_0 \approx 53 \text{ km} \cdot \text{sec}^{-1} \cdot \text{Mpc}^{-1}$ for the Hubble constant, and 3.- We obtain a bad fit to data with a $\chi^2_{\min} \approx 532$ ($\chi^2_{\text{d.o.f.}} \approx 2.92$). These results indicate that this model is viable just if the bulk viscosity is triggered in recent times.

We present a flat cosmological model which component is a pressureless fluid made of baryon and dark matter components with a bulk viscosity coefficient ζ_m proportional to the Hubble parameter, i.e., $\zeta_m = \zeta H$, where $\zeta = \text{constant}$ and H is the Hubble parameter, to explain the present accelerated expansion of the universe. The subscript “m” stands for *matter* with bulk viscosity. A bulk viscosity coefficient can produce a positive term in the second Friedmann equation that induces an acceleration [1, 2] (i.e., $\ddot{a} \geq 0$, with a denoting the scale factor and the dots derivatives with respect to the cosmic time). Similar models or analysis have been proposed also in [2, 3, 4, 5, 6].

A bulk viscous matter-dominated universe The energy-momentum tensor of a fluid composed by *only* matter (hereinafter we call *matter* to baryon

and dark matter components together) with bulk viscosity ζ_m is defined by $T_{\mu\nu} = \rho_m u_\mu u_\nu + (g_{\mu\nu} + u_\mu u_\nu) P_m^*$, where $P_m^* \equiv P_m - 3\zeta_m (\dot{a}/a)$ (see e.g. [1]). Here P_m is the pressure of the matter fluid and the function a the scale factor. The four-velocity vector u_μ is that of a comoving observer that measures the pressure P_m^* and the density ρ_m of the matter fluid. Assuming pressureless matter $P_m = 0$ implies $P_m^* = -3\zeta_m (\dot{a}/a)$.

On the other hand, the conservation of energy has the form $a(d\rho_m/da) = 3(3\zeta_m H - \rho_m)$, where $H \equiv \dot{a}/a$ is the Hubble parameter. Using the ansatz $\zeta_m = \zeta H$ for the parametrization of the bulk viscosity¹ and the first Friedmann equation $H^2 = (8\pi G/3) \rho_m$ (assuming a spatially flat geometry for the universe), the conservation equation becomes $a(d\rho_m/da) = 3\rho_m [(8\pi G/3) \zeta - 1]$. The solution in terms of the redshift² ‘ z ’ is $\rho_m(z) = \rho_m^0 (1+z)^{3-\tilde{\zeta}}$, where we have defined the dimensionless bulk viscous coefficient $\tilde{\zeta} \equiv 8\pi G \zeta$ and ρ_m^0 is the value of the density of matter evaluated today. We substitute this solution into the first Friedmann equation so it becomes $H^2 = (8\pi G/3) \rho_m^0 (1+z)^{3-\tilde{\zeta}}$. Dividing the last expression by the *critical density* today $\rho_{\text{crit}}^0 \equiv 3H_0^2/8\pi G$ we obtain

$$H(z) = H_0(1+z)^{\frac{1}{2}(3-\tilde{\zeta})} \quad (1)$$

where H_0 is the Hubble constant. In this model the matter density parameter is $\Omega_m^0 \equiv \rho_m^0/\rho_{\text{crit}}^0 = 1$ because the matter is the only component of the universe.

Cosmological tests We constrain the possible values of the bulk viscosity $\tilde{\zeta}$ using the following cosmological tests: the Gold 2006 SNe Ia data sample [7] composed by 182 SNe Ia, the Cosmic Microwave Background (CMB) shift parameter ‘ R ’ from the three-year WMAP observations [8], the Baryon Acoustic Oscillation (BAO) peak ‘ A ’ from the Sloan Digital Sky Survey (SDSS) [9] and the Second Law of Thermodynamics (SLT) [1].

For the SNe Ia test it is defined the observational luminosity distance [10, 11] in a flat cosmology as $d_L = c(1+z)H_0^{-1} \int_0^z E(z')^{-1} dz'$, where $E(z) \equiv H(z)/H_0$ and c the speed of light. The *theoretical distance moduli* for the i -th supernovae with redshift z_i is $\mu(z_i) = 5 \log_{10}[d_L(z_i)/\text{Mpc}] + 25$. The statistical function χ_{SNe}^2 becomes $\chi_{\text{SNe}}^2(\tilde{\zeta}, H_0) \equiv \sum_{k=1}^{182} [\mu(z_k, \tilde{\zeta}, H_0) - \mu_k]^2 / \sigma_k^2$, where μ_k is the *observed* distance moduli for the k -th supernovae and σ_k^2 is the variance of the measurement.

The CMB shift parameter R is defined as $R \equiv \sqrt{\Omega_m^0} \int_0^{Z_{\text{CMB}}} E(z')^{-1} dz'$, where $Z_{\text{CMB}} = 1089$ is the redshift of recombination [8, 12]. The observed value of the shift parameter R is reported to be $R_{\text{obs}} = 1.70 \pm 0.03$ [8].

From the SDSS data, we use the baryon acoustic oscillation (BAO) peak A defined as $A \equiv \sqrt{\Omega_m^0} E(z_1)^{-1/3} [z_1^{-1} \int_0^{z_1} E(z')^{-1} dz']^{2/3}$, where $z_1 = 0.35$ [9]. The observed value of SDSS-BAO peak is $A_{\text{obs}} = 0.469 \pm 0.017$ [9].

¹The explicit form of the bulk viscosity is assumed a prior. See for instance [2, 3, 5, 6].

²The superscript ‘‘0’’ (zero) represents a quantity evaluated today and the relation between the redshift z and the scale factor a is $(1+z) = 1/a$.




Figure 1: Confidence intervals for the parameters $(\tilde{\zeta}, H_0)$ for a spatially flat bulk viscous matter-dominated universe. We show the confidence intervals of 68.3%(1 σ), 95.4%(3 σ) and 99.73%(5 σ). The constraints are derived from the Gold 2006 SNe Ia data sample alone (left), and the joint Gold 2006 + CMB + BAO cosmological tests (right). H_0 is in units of $\text{km} \cdot \text{sec}^{-1} \cdot \text{Mpc}^{-1}$ and $\tilde{\zeta}$ is dimensionless ($\tilde{\zeta} \equiv 8\pi G\zeta$).

We construct the total χ^2 function as $\chi^2_{\text{total}} \equiv \chi^2_{\text{SNe}} + \chi^2_{\text{CMB}} + \chi^2_{\text{BAO}}$, where χ^2_{SNe} was defined above, $\chi^2_{\text{CMB}} \equiv [(R - R_{\text{obs}})/\sigma_R]^2$ and $\chi^2_{\text{BAO}} \equiv [(A - A_{\text{obs}})/\sigma_A]^2$.

The law of generation of *local* entropy in the space-time is found to be $T \nabla_\nu s^\nu = \zeta_m \nabla_\nu u^\nu = 3H\zeta_m$ [1], where T is the temperature, $\nabla_\nu s^\nu$ is the rate at which entropy is being generated in a unit volume. The second law of thermodynamics can be written as $T \nabla_\nu s^\nu \geq 0$ which implies that $3H\zeta_m \geq 0$. Since H is positive in an expanding universe then $\zeta_m = \zeta H$ has to be positive in order to preserve the validity of the second law of thermodynamics. Thus, in the present model a necessary condition is $\tilde{\zeta} \geq 0$.

Constraining the bulk viscous cosmological model We compute the *best estimated values* and “the goodness-of-fit” of $\tilde{\zeta}$ and H_0 to the data through χ^2 -minimization, using SNe Ia test alone, and also considering the joint SNe Ia, CMB and BAO tests together. Then we compute the confidence intervals for $(\tilde{\zeta}, H_0)$ to constrain their possible values. We obtain as best estimates using only the SNe Ia test: $\tilde{\zeta} = 1.43 \pm 0.12$ and $H_0 = 61.758 \pm 0.86$, with a $\chi^2_{\text{min}} = 165.135$ ($\chi^2_{\text{d.o.f.}} = 0.917$), where H_0 is in units of $\text{km} \cdot \text{sec}^{-1} \cdot \text{Mpc}^{-1}$. On the other hand, using the SNe Ia, CMB and BAO tests together we obtain $\tilde{\zeta} = -0.1451 \pm 0.02$ and $H_0 = 53.425 \pm 0.4$, with a $\chi^2_{\text{min}} = 532.056$ ($\chi^2_{\text{d.o.f.}} = 2.92$). It can be seen that when we consider only the SNe Ia test we obtain $\tilde{\zeta} \geq 0$ with a 99.7% confidence level and a reasonable value for the Hubble constant in the interval $58.9 \leq H_0 \leq 64.8$ as well as for $\chi^2_{\text{d.o.f.}} = 0.917$. However, when the same analysis is performed using the three cosmological tests together (SNe Ia, CMB




Figure 2: Confidence intervals for the parameters $(\tilde{\zeta}, H_0)$ for a spatially flat bulk viscous matter-dominated universe. We show the confidence intervals of 68.3%(1 σ), 95.4%(3 σ) and 99.73%(5 σ). The constraints are derived from the Gold 2006 SNe Ia data sample alone and the joint Gold 2006 + CMB + BAO cosmological tests. H_0 is in units of $\text{km} \cdot \text{sec}^{-1} \cdot \text{Mpc}^{-1}$ and $\tilde{\zeta}$ is dimensionless ($\tilde{\zeta} \equiv 8\pi G\zeta$).

and BAO) we obtain negative values of $\tilde{\zeta}$ with at least 99.7 % confidence level that disagree with the second law of thermodynamics, a not so good estimation for H_0 of $52.1 \leq H_0 \leq 54.8$, and a bad $\chi^2_{\text{d.o.f.}} = 2.92$. These results are illustrated in figures 1 and 2.

Therefore, we conclude of this model that:

- In order to have a viable model for explaining the observed accelerated expansion of the universe, the bulk viscosity of the fluid with the ansatz considered in the present work should be triggered in recent times ($z \lesssim 2$).
- Since the CMB and BAO tests have information for very large redshifts ($z \gg 2$) and due to the fact of that this model has a bad fit and violate the second law of thermodynamics when we test it using the joint SNe Ia, CMB and BAO tests then we conclude that this is not viable if it is extrapolated for early times of the universe as suggested in [5].

Acknowledgments This work is partly supported by grants CIC-UMSNH 4.8, PROMEP UMSNH-CA-22, SNI-20733 and COECYT.

References

- [1] C. W. Misner, K. S. Thorne and J. A. Wheeler, **Gravitation**, W. H. Freeman and Company, p 567, (1973).
- [2] J. C. Fabris, S. V. B. Goncalves and R. de Sa Ribeiro, *Gen. Rel. Grav* **38**, 495–506, (2006).
- [3] A. Avelino, U. Nucamendi and F. S. Guzman, in *XI Mexican Workshop on Particles and Fields. AIP Conf. Proc*, AIP **1026**, 300–302, arXiv:0801.1686 [gr-qc], (2008).
- [4] Arturo Avelino and Ulises Nucamendi, *in preparation*, (2008).
- [5] R. Colistete et al., *Phys. Rev. D* **76**, 103516, 1–13, (2007).
- [6] Jie Ren and Xin-He Meng, *Phys. Lett. B* **633**, 1–8, (2006).
- [7] Adam G. Riess et al., *Astrophysical journal* **659**, 98–121, (2007).
- [8] Y. Wang and P. Mukherjee, *Astrophysical Journal* **650**, 1–6, (2006).
- [9] D. J. Eisenstein et al., (SDSS Collaboration) *Astrophysical Journal* **633**, 560–574, (2005).
- [10] M. Turner and A. G. Riess, *Astrophysical journal* **569**, 18, (2002).
- [11] A. G. Riess et al., *Astrophysical journal* **607**, 665–687, (2004).
- [12] A. Melchiorri et al., *Phys. Rev. D* **68**, 043509, (2003).